

# Minimal Retentive Sets in Tournaments

Felix Brandt

Markus Brill

Felix Fischer

Paul Harrenstein

Institut für Informatik  
Universität München  
80538 München, Germany  
{brandtf,brill,fischerf,harrenst}@tcs.ifi.lmu.de

## ABSTRACT

Many problems in multiagent decision making can be addressed using *tournament solutions*, i.e., functions that associate with each complete and asymmetric relation on a set of alternatives a non-empty subset of the alternatives. For a given tournament solution  $S$ , Schwartz calls a set of alternatives  $S$ -retentive if it satisfies a natural stability criterion with respect to  $S$ . He then recursively defines the *tournament equilibrium set (TEQ)* as  $\mathring{TEQ}$ , the union of all inclusion-minimal  $TEQ$ -retentive sets. Due to its unwieldy recursive definition, precious little is known about  $TEQ$ . Assuming a well-known conjecture about  $TEQ$ , we show that most desirable properties of tournament solutions are inherited from  $S$  to  $\mathring{S}$ . We thus obtain an infinite hierarchy of efficiently computable tournament solutions that “approximate”  $TEQ$  (which is computationally intractable) while maintaining most of its desirable properties. This hierarchy contains well-known tournament solutions such as the top cycle ( $TC$ ) and the minimal covering set ( $MC$ ). We prove a weaker version of the conjecture mentioned above, which establishes  $\mathring{TC}$  as an attractive new tournament solution.

## Categories and Subject Descriptors

I.2.11 [Distributed Artificial Intelligence]: Multiagent Systems; J.4 [Computer Applications]: Social and Behavioral Sciences—*Economics*

## General Terms

Theory, Economics

## Keywords

Social Choice Theory, Tournament Solutions, Retentiveness, Tournament Equilibrium Set

## 1. INTRODUCTION

A tournament solution is a function that associates with each complete and asymmetric relation on a set of alternatives a non-empty subset of the alternatives. Tournament solutions thus provide a framework in which many problems in

**Cite as:** Minimal Retentive Sets in Tournaments, Felix Brandt, Markus Brill, Felix Fischer, and Paul Harrenstein, *Proc. of 9th Int. Conf. on Autonomous Agents and Multiagent Systems (AAMAS 2010)*, van der Hoek, Kaminka, Luck and Sen (eds.), May, 10–14, 2010, Toronto, Canada, pp. XXX-XXX.

Copyright © 2010, International Foundation for Autonomous Agents and Multiagent Systems (www.ifaamas.org). All rights reserved.

multiagent decision making can be addressed. For instance, they play an important role in social choice theory, where the binary relation is typically defined via pairwise majority voting [22, 21]. Other application areas include multi-criteria decision analysis [2, 3], zero-sum games [14, 19, 10], coalition formation [6], and argumentation theory [11, 12].

Examples of well-studied tournament solutions are the minimal covering set, the Banks set, and the Slater set [21]. Recent years have witnessed an increasing interest in these concepts by the multiagent systems and theoretical computer science communities, particularly with respect to their computational complexity. For example, it was shown that the minimal covering set of a tournament can be computed in polynomial time [5], whereas computing the Banks set and the Slater set is computationally intractable [25, 1, 9].

The *tournament equilibrium set (TEQ)* introduced by Schwartz [24] ranks among the most intriguing, but also among the most enigmatic, tournament solutions. For a given tournament solution  $S$ , Schwartz calls a set of alternatives  $S$ -retentive if it satisfies a natural stability criterion with respect to  $S$ . He then recursively defines  $TEQ$  as  $\mathring{TEQ}$ , the union of all inclusion-minimal  $TEQ$ -retentive sets. Unfortunately, and somewhat surprisingly, it is unknown whether  $TEQ$  satisfies any of several important properties proposed in the literature on tournament solutions, namely *monotonicity*, *independence of unchosen alternatives*, and the *weak superset property*. However, Laffond et al. [18] and Houy [16, 17] have shown that  $TEQ$  satisfies one of these properties if and only if it satisfies all of them. They moreover showed that  $TEQ$  satisfying any of the properties is equivalent to the statement that every tournament contains a *unique* minimal  $TEQ$ -retentive set. This statement had already been conjectured by Schwartz [24] and also implies that  $TEQ$  is strictly contained in the minimal covering set. Apart from these implications, the only known facts about  $TEQ$  are that it is contained in the Banks set [24], satisfies composition-consistency [20], and is NP-hard to compute [8].

In this paper, we generally study tournament solutions that are defined via Schwartz’s notion of retentiveness, i.e., we consider  $\mathring{S}$  for any given tournament solution  $S$ . For tournament solutions  $S$  that always admit a unique minimal  $S$ -retentive set, we show that most desirable properties are inherited from  $S$  to  $\mathring{S}$  (and also from  $\mathring{S}$  to  $S$ ). Composition-consistency is a notable exception as we prove that  $TEQ$  is the *only* composition-consistent tournament solution defined via retentiveness.

Starting with the trivial tournament solution that always returns all alternatives, one can define an infinite sequence

of tournament solutions  $(S_1, S_2, \dots)$  such that  $S_{i+1} = \mathring{S}_i$ . Assuming Schwartz's conjecture, we show that these tournament solutions are strictly contained in each other, strictly contain  $TEQ$ , and share most of the desirable properties of  $TEQ$ . The sequence converges (in a well-defined way) to  $TEQ$  and yields an infinite sequence of weaker versions of Schwartz's conjecture. The first statement of this sequence was shown by Good [15], and we conclude the paper by proving the second one.

## 2. PRELIMINARIES

The central question in the theory of tournament solutions is how to extend choices in sets consisting of only two elements to larger sets. Choices over pairs of alternatives can conveniently be represented by a tournament, i.e., a binary relation on the entire set of alternatives. Tournament solutions then advocate different views on how to choose from arbitrary subsets of alternatives based on these pairwise comparisons (see [21], for an excellent overview of tournament solutions and their properties).

### 2.1 Tournaments

Let  $X$  be a universe of alternatives, and assume for notational convenience that  $\mathbb{N} \subseteq X$ . The set of all *finite* subsets of  $X$  will be denoted by  $\mathcal{F}_0(X)$ , the set of all *non-empty* finite subsets of  $X$  by  $\mathcal{F}(X)$ . A (finite) *tournament*  $T$  is a pair  $(A, \succ)$ , where  $A \in \mathcal{F}(X)$  and  $\succ$  is an asymmetric and complete (and thus irreflexive) binary relation on  $X$ , usually referred to as the *dominance relation*.<sup>1</sup> Intuitively,  $a \succ b$  signifies that alternative  $a$  is preferable to alternative  $b$ . The dominance relation can be extended to sets of alternatives by writing  $A \succ B$  when  $a \succ b$  for all  $a \in A$  and  $b \in B$ . We further write  $\mathcal{D}(X)$  for the set of all dominance relations on  $X$ , and  $\mathcal{T}(X) = \mathcal{F}(X) \times \mathcal{D}(X)$  for the set of all tournaments on  $X$ .

For a tournament  $(A, \succ)$  and an element  $a \in A$ , we denote by  $D_\succ(a)$  the *dominion* of  $a$ , i.e.,

$$D_\succ(a) = \{b \in A : a \succ b\},$$

and by  $\bar{D}_\succ(a)$  the *dominators* of  $a$ , i.e.,

$$\bar{D}_\succ(a) = \{b \in A : b \succ a\}.$$

Whenever the dominance relation  $\succ$  is known from the context, the subscript will be omitted to improve readability. For a tournament  $T = (A, \succ)$  and a subset  $B \subseteq A$  of alternatives, we further write  $T|_B = (B, \{(a, b) \in B \times B : a \succ b\})$  for the restriction of  $T$  to  $B$ .

The *order* of a tournament  $T = (A, \succ)$  refers to the cardinality of  $A$ . A *tournament isomorphism* of two tournaments  $T = (A, \succ)$  and  $T' = (A', \succ')$  is a bijection  $\pi : A \rightarrow A'$  such that  $a \succ b$  if and only if  $\pi(a) \succ' \pi(b)$ .

### 2.2 Components and Decompositions

An important structural notion in the context of tournaments is that of a *component*. A component is a subset of alternatives that bear the same relationship to all alternatives not in the set.

<sup>1</sup>This definition slightly diverges from the common graph-theoretic definition where  $\succ$  is defined on  $A$  rather than  $X$ . It however facilitates the sound definition of tournament solutions.

DEFINITION 1. Let  $T = (A, \succ)$  be a tournament. A *non-empty subset*  $B$  of  $A$  is a *component* of  $T$  if for all  $a \in A \setminus B$ , either  $B \succ a$  or  $a \succ B$ . A *decomposition* of  $T$  is a set of disjoint components  $\{B_1, \dots, B_k\}$  of  $T$  such that  $A = \bigcup_{i=1}^k B_i$ .

For a given tournament  $\tilde{T}$ , a new tournament can be constructed by replacing each alternative with a component.

DEFINITION 2. Let  $B_1, \dots, B_k \subseteq X$  be disjoint sets and  $\tilde{T} = (\{1, \dots, k\}, \tilde{\succ})$ ,  $T_1 = (B_1, \succ_1)$ ,  $\dots$ ,  $T_k = (B_k, \succ_k)$  tournaments. The *product* of  $T_1, \dots, T_k$  with respect to  $\tilde{T}$ , denoted by  $\Pi(\tilde{T}, T_1, \dots, T_k)$ , is the tournament  $(A, \succ)$  such that  $A = \bigcup_{i=1}^k B_i$  and for all  $b_i \in B_i, b_j \in B_j$ ,

$$b_i \succ b_j \quad \text{if only if} \quad i = j \text{ and } b_i \succ_i b_j, \text{ or } i \neq j \text{ and } i \tilde{\succ} j.$$

### 2.3 Tournament Solutions

Consider the *maximum* function  $\max : \mathcal{T}(X) \rightarrow \mathcal{F}_0(X)$  given by  $\max((A, \succ)) = \{a \in A : a \succ b \text{ for all } b \in A \setminus \{a\}\}$ . Due to the asymmetry of the dominance relation, this function returns at most one alternative in any tournament. Moreover, maximal—i.e., undominated—and maximum elements coincide. In social choice theory, the maximum of a majority tournament is commonly referred to as the *Condorcet winner*.

Since the dominance relation may contain cycles and thus fail to have a maximal element, a variety of concepts have been suggested to take over the role of singling out the “best” alternatives of a tournament. Formally, a *tournament solution*  $S$  is defined as a function that associates with each tournament  $T = (A, \succ)$  a non-empty subset  $S(T)$  of  $A$ . Following Laslier [21], we require a tournament solution to be independent of alternatives outside the tournament, invariant under tournament isomorphisms, and to select the maximal element whenever it exists.

DEFINITION 3. A *tournament solution* is a function  $S : \mathcal{T}(X) \rightarrow \mathcal{F}(X)$  such that

- (i)  $S(T) = S(T')$  for all tournaments  $T = (A, \succ)$  and  $T' = (A, \succ')$  such that  $T|_A = T'|_A$ ;
- (ii)  $S((\pi(A), \succ')) = \pi(S((A, \succ)))$  for all tournaments  $(A, \succ)$ ,  $(A', \succ')$ , and every tournament isomorphism  $\pi : A \rightarrow A'$  of  $(A, \succ)$  and  $(A', \succ')$ ; and
- (iii)  $\max(T) \subseteq S(T) \subseteq A$  for all tournaments  $T = (A, \succ)$ .

Laslier [21] is slightly more stringent here as he requires the maximum to be the only element in  $S(T)$  whenever it exists. We will call a tournament solution *proper* if it satisfies this additional requirement.

The conditions of Definition 3 are trivially satisfied if one invariably selects the set of all alternatives. The corresponding tournament solution  $TRIV$  is obtained by letting  $TRIV((A, \succ)) = A$  for every tournament  $(A, \succ)$ . Among the tournament solutions considered in this paper,  $TRIV$  is the only one that is not proper. The *top cycle*  $TC(T)$  of a tournament  $T = (A, \succ)$  is defined as the smallest set  $B \subseteq A$  such that  $B \succ A \setminus B$ . Uniqueness of such a set is straightforward and was first shown by Good [15].

For two tournament solutions  $S$  and  $S'$ , we write  $S' \subseteq S$ , and say that  $S'$  is a *refinement* of  $S$ , if  $S'(T) \subseteq S(T)$  for all tournaments  $T$ . To avoid cluttered notation, we write  $S(A, \succ)$  instead of  $S((A, \succ))$  for a tournament  $T = (A, \succ)$ .

Furthermore, we frequently write  $S(B)$  instead of  $S(T|_B)$  for a subset  $B \subseteq A$  of alternatives, if the dominance relation  $\succ$  is known from the context.

## 2.4 Retentive Sets

Motivated by cooperative majority voting, Schwartz [24] introduced a tournament solution based on a notion he calls *retentiveness*. The intuition underlying retentive sets is that alternative  $a$  is only “properly” dominated by alternative  $b$  if  $b$  is chosen among  $a$ ’s dominators by some underlying tournament solution  $S$ . A set of alternatives is then called  $S$ -retentive if none of its elements is properly dominated by some alternative outside the set with respect to  $S$ .

**DEFINITION 4.** *Let  $S$  be a tournament solution and  $T = (A, \succ)$  a tournament. Then,  $B \subseteq A$  is  $S$ -retentive in  $T$  if  $B \neq \emptyset$  and  $S(\overline{D}(b)) \subseteq B$  for all  $b \in B$  such that  $\overline{D}(b) \neq \emptyset$ . The set of  $S$ -retentive sets for a given tournament  $T = (A, \succ)$  will be denoted by  $\mathcal{R}_S(T)$ , i.e.,  $\mathcal{R}_S(T) = \{B \subseteq A : B \text{ is } S\text{-retentive in } T\}$ .*

Fix an arbitrary tournament solution  $S$ . Since the set  $A$  of all alternatives is trivially  $S$ -retentive in  $(A, \succ)$ ,  $S$ -retentive sets are guaranteed to exist. If a Condorcet winner exists, it must clearly be contained in any  $S$ -retentive set. The union of all (inclusion-)minimal  $S$ -retentive sets thus defines a tournament solution.

**DEFINITION 5.** *Let  $S$  be a tournament solution. Then, the tournament solution  $\hat{S}$  is given by*

$$\hat{S}(T) = \bigcup_{\subseteq} \min(\mathcal{R}_S(T)).$$

Consider for example the tournament solution  $TRIV$ , which always selects the set of all alternatives. It is easily verified that there always exists a *unique* minimal  $TRIV$ -retentive set, and that in fact  $\hat{TRIV} = TC$ .

For a tournament solution  $S$ , we say that  $\mathcal{R}_S$  is *pairwise intersecting* if for each tournament  $T$  and for all sets  $B, C \in \mathcal{R}_S(T)$ ,  $B \cap C \neq \emptyset$ . Observe that the non-empty intersection of two  $S$ -retentive sets is itself  $S$ -retentive. We thus have the following.

**PROPOSITION 1.** *For every tournament solution  $S$ ,  $\mathcal{R}_S$  admits a unique minimal element if and only if  $\mathcal{R}_S$  is pairwise intersecting.*

Schwartz introduced retentiveness in order to recursively define the *tournament equilibrium set (TEQ)* as the union of minimal  $TEQ$ -retentive sets. This recursion is well-defined because the order of the dominator set of any alternative is strictly smaller than the order of the original tournament.

**DEFINITION 6** (Schwartz [24]). *The tournament equilibrium set (TEQ) is defined recursively as  $TEQ = \hat{TEQ}$ .*

In other words,  $TEQ$  is the unique fixed point of the  $\circ$ -operator. Schwartz conjectured that every tournament admits a *unique* minimal  $TEQ$ -retentive set.

**CONJECTURE 1** (Schwartz [24]).  *$\mathcal{R}_{TEQ}$  is pairwise intersecting.*

Despite several attempts to prove or disprove this statement (e.g., [18, 16]), it has remained a conjecture. A recent computer analysis failed to find a counter-example in all

tournaments of order 12 or less and a fairly large number of random tournaments [8].

It turns out that the existence of a *unique* minimal  $S$ -retentive set is quintessential for showing that  $\hat{S}$  satisfies several important properties to be defined the next section.

## 2.5 Properties of Tournament Solutions

In order to compare tournament solutions with each other, a number of desirable properties for tournament solutions have been identified. In this section, we will review six of the most common properties.<sup>2</sup> Moulin [23], in a more general context, distinguishes between *monotonicity* and *independence* conditions, where a monotonicity condition describes the positive association of the solution with some parameter, and an independence condition characterizes the invariance of the solution under the modification of some parameter. Properties of tournament solutions can further be distinguished depending on whether they are defined via the dominance relation or via set inclusion.

We first consider a monotonicity and an independence property defined in terms of the dominance relation. A tournament solution is called *monotonic* if a chosen alternative remains in the choice set when extending its dominion and leaving everything else unchanged.

**DEFINITION 7.** *A tournament solution  $S$  satisfies monotonicity (MON) if  $a \in S(T)$  implies  $a \in S(T')$  for all tournaments  $T = (A, \succ)$ ,  $T' = (A, \succ')$ , and  $a \in A$  such that  $T|_{A \setminus \{a\}} = T'|_{A \setminus \{a\}}$  and  $D_{\succ}(a) \subseteq D_{\succ'}(a)$ .*

A solution satisfies *independence of unchosen alternatives* if the choice set is invariant under any modification of the dominance relation between unchosen alternatives.

**DEFINITION 8.** *A tournament solution  $S$  is independent of unchosen alternatives (IUA) if  $S(T) = S(T')$  for all tournaments  $T = (A, \succ)$  and  $T' = (A, \succ')$  such that  $T|_{S(T) \cup \{a\}} = T'|_{S(T) \cup \{a\}}$  for all  $a \in A$ .*

With respect to set inclusion, we consider a monotonicity property to be called the *weak superset property* and an independence property known as the *strong superset property*. A tournament solution satisfies the weak superset property if an unchosen alternative remains unchosen when other unchosen alternatives are removed.


**DEFINITION 9.** *A tournament solution  $S$  satisfies the weak superset property (WSP) if  $S(B) \subseteq S(A)$  for all tournaments  $T = (A, \succ)$  and  $B \subseteq A$  such that  $S(A) \subseteq B$ .*

The strong superset property requires that a choice set is invariant under the removal of alternatives not in the choice set.

**DEFINITION 10.** *A tournament solution  $S$  satisfies the strong superset property (SSP) if  $S(B) = S(A)$  for all tournaments  $T = (A, \succ)$  and  $B \subseteq A$  such that  $S(A) \subseteq B$ .*

The four properties defined above (MON, IUA, WSP, and SSP) will be called *basic* properties of tournament solutions.

<sup>2</sup>Our terminology slightly differs from the one by Laslier [21] and others. *Independence of unchosen alternatives* is also called *independence of the losers* or *independence of non-winners*. The *weak superset property* has been referred to as  $\epsilon^+$  or as the *Aizerman property*.



**Figure 1:** Tournament  $C(T, I_a, I_b)$  for a given tournament  $T$ . The gray circle represents a component isomorphic to the original tournament  $T$ . An edge incident to a component signifies that there is an edge of the same direction incident to each alternative in the component.

Observe that SSP implies WSP. Furthermore, the conjunction of MON and SSP implies IUA. It is therefore sufficient to show MON and SSP in order to prove that a tournament solution satisfies all basic properties. An additional properties considered in this paper is *composition-consistency*. A tournament solution is composition-consistent if it chooses the “best” alternatives from the “best” components.

**DEFINITION 11.** A tournament solution  $S$  is composition-consistent (COM) if for all tournaments  $T, T_1, \dots, T_k$ , and  $\tilde{T}$  such that  $T = \Pi(\tilde{T}, T_1, \dots, T_k)$ ,  $S(T) = \bigcup_{i \in S(\tilde{T})} S(T_i)$ .

The properties defined in this section are not easily satisfied by discriminative tournament solutions. While *TRIV* trivially satisfies all of the properties, the Slater set only satisfies MON and the Banks set only satisfies MON, WSP, and COM. The minimal covering set satisfies all of the properties. The same holds for *TEQ* if Conjecture 1 is correct.

### 3. INHERITANCE OF PROPERTIES

In this section, we investigate which of the properties defined in the previous section are inherited from  $S$  to  $\hat{S}$  or from  $\hat{S}$  to  $S$ .

We begin by looking at a particular type of decomposable tournament that will be useful in the following. Let  $C_3$  denote the tournament  $C_3 = (\{1, 2, 3\}, \succ)$  with  $1 \succ 2 \succ 3 \succ 1$ , and write  $I_a$  for the unique tournament on  $\{a\}$ . For three tournaments  $T_1, T_2$ , and  $T_3$ , let  $C(T_1, T_2, T_3) = \Pi(C_3; T_1, T_2, T_3)$ . The structure of  $C(T, I_a, I_b)$  for a given tournament  $T$  is illustrated in Figure 1. We have the following lemma.

**LEMMA 1.** Let  $S$  be a proper tournament solution. Then, for each tournament  $T$  on  $A$  and all  $a, b \notin A$ ,

$$\hat{S}(C(T, I_a, I_b)) = \{a, b\} \cup S(T).$$

**PROOF.** Let  $B = \hat{S}(C(T, I_a, I_b))$ , and observe that  $B \cap A \neq \emptyset$ , because neither  $\{a, b\}$  nor any subset of it is  $S$ -retentive. Since  $a$  is the Condorcet winner in  $\bar{D}(b) = \{a\}$  and  $b$  is the Condorcet winner in  $\bar{D}(a) = \{b\}$ , by  $S$ -retentiveness of  $B$  we must have that  $a \in B$  and  $b \in B$ . Also by retentiveness of  $B$ , we have  $S(\bar{D}(a)) = S(T) \subseteq B$ . We have thus shown that every  $S$ -retentive set must contain  $\{a, b\} \cup S(T)$ , and that  $\{a, b\} \cup S(T)$  is itself  $S$ -retentive.  $\square$

We are now ready to show that many desirable properties (including efficient computability) are inherited from  $\hat{S}$  to  $S$  and from  $S$  to  $\hat{S}$ .

**THEOREM 1.** Let  $S$  be a proper tournament solution. Then the following holds:

- (i) Each of the four basic properties is satisfied by  $S$  if and only if it is satisfied by  $\hat{S}$ .
- (ii) If  $\mathcal{R}_S$  is pairwise intersecting, each of the following properties is satisfied by  $\hat{S}$  if and only if it is satisfied by  $S$ : (MON  $\wedge$  SSP), SSP, WSP, IUA.
- (iii)  $\hat{S}$  is efficiently computable if and only if  $S$  is efficiently computable.

**PROOF.** For (i), we show the following: If  $S$  violates one of the four basic properties MON, SSP, WSP, or IUA, then  $\hat{S}$  violates the same property. Observe that for each of these properties, the fact that  $S$  violates the property can be witnessed by a pair of tournaments  $T_1 = (A_1, \succ_1)$  and  $T_2 = (A_2, \succ_2)$ : In the case of SSP (resp., WSP),  $S(T_1) \subseteq A_2 \subset A_1, T_2|_{A_2} = T_1|_{A_2}$ , and  $S(T_2) \neq S(T_1)$  (resp.,  $S(T_2) \subsetneq S(T_1)$ ). In the case of MON and IUA,  $A_2 = A_1$  and the only difference between the dominance relations is that  $D_{\succ_2}(a) = D_{\succ_1}(a) \cup \{b\}$  for some alternatives  $a, b \in A_1$ . For MON,  $a \in S(T_1)$  and  $a \notin S(T_2)$ ; for IUA,  $a, b \notin S(T_1)$  and  $S(T_1) \neq S(T_2)$ .

The pair  $(T_1, T_2)$  is called a counterexample. We go on to show how a counterexample for  $S$  can be transformed into a counterexample for  $\hat{S}$ . For  $a, b \notin A_1$ , define  $T'_1 = C(T_1, I_a, I_b)$  and  $T'_2 = C(T_2, I_a, I_b)$ . Lemma 1 implies that  $\hat{S}(T'_1) = \{a, b\} \cup S(T_1)$  and  $\hat{S}(T'_2) = \{a, b\} \cup S(T_2)$ . Hence, the pair  $(T'_1, T'_2)$  constitutes a counterexample for  $\hat{S}$ .

For (ii), assume that  $\mathcal{R}_S$  is directed. We need to show that each of the properties (MON  $\wedge$  SSP), SSP, WSP, and IUA is satisfied by  $S$  if and only if it is satisfied by  $\hat{S}$ . The direction from left to right follows from (i). We now show that the properties are inherited from  $S$  to  $\hat{S}$ .

Assume  $S$  satisfies SSP. Let  $T$  be a tournament on  $A$ , and consider an alternative  $x \in A \setminus \hat{S}(T)$ . Let  $T' = T|_{A \setminus \{x\}}$ . Since  $\mathcal{R}_S$  is pairwise intersecting, it suffices to show that for all  $a \in \hat{S}(T)$ ,  $S(\bar{D}_A(a)) = S(\bar{D}_{A \setminus \{x\}}(a))$ . To this end, consider an arbitrary  $a \in \hat{S}(T)$ . If  $x \notin \bar{D}_A(a)$ , then obviously  $\bar{D}_A(a) = \bar{D}_{A \setminus \{x\}}(a)$  and thus  $S(\bar{D}_A(a)) = S(\bar{D}_{A \setminus \{x\}}(a))$ . Assume on the other hand that  $x \in \bar{D}_A(a)$ . Since  $x \notin \hat{S}(T)$  and  $a \in \hat{S}(T)$ , it follows that  $x \notin S(\bar{D}_A(a))$ . Now, since  $S$  satisfies SSP, we obtain  $S(\bar{D}_A(a)) = S(\bar{D}_{A \setminus \{x\}}(a))$  as desired.

Assume that  $S$  satisfies WSP. Let  $T$  be a tournament on  $A$ , and consider an alternative  $x \in A \setminus \hat{S}(T)$ . Since  $\mathcal{R}_S$  is pairwise intersecting, it suffices to show that  $\hat{S}(T)$  is also  $S$ -retentive in  $T|_{A \setminus \{x\}}$ . To this end, consider an arbitrary  $a \in \hat{S}(T)$ . Since  $S$  satisfies WSP, we have that  $S(\bar{D}_{A \setminus \{x\}}(a)) \subseteq S(\bar{D}_A(a))$ . Furthermore, by  $S$ -retentiveness of  $\hat{S}(T)$ ,  $S(\bar{D}_A(a)) \subseteq \hat{S}(T)$  and thus  $S(\bar{D}_{A \setminus \{x\}}(a)) \subseteq \hat{S}(T)$ .

Assume that  $S$  satisfies IUA. Let  $T$  and  $T'$  be two tournaments on  $A$ , and consider  $x, y \in A \setminus \hat{S}(T)$  such that  $T|_{A \setminus \{x, y\}} = T'|_{A \setminus \{x, y\}}$  and  $x \succ y$  if and only if  $y \succ' x$ . Since  $\mathcal{R}_S$  is pairwise intersecting, it suffices to show that for all  $a \in \hat{S}(T)$ ,

$$S(T|_{\bar{D}(a)}) = S(T'|_{\bar{D}(a)}),$$

which directly implies that  $S(T)$  is also  $S$ -retentive in  $T'$ . To this end, consider an arbitrary  $a \in \hat{S}(T)$ . By assumption,  $a \neq x$  and  $a \neq y$ . First consider the case when both  $x \in$

$\overline{D}_T(a)$  and  $y \in \overline{D}_T(a)$ . Then,  $\overline{D}_T(a) = \overline{D}_{T'}(a)$  and, by  $S$ -retentiveness of  $S(T)$ ,  $x, y \notin S(T|_{\overline{D}_T(a)})$ . Since  $S$  satisfies IUA,  $S(T|_{\overline{D}_T(a)}) = S(T|_{\overline{D}_{T'}(a)})$  as required. Now consider the case when  $x \notin \overline{D}_T(a)$  or  $y \notin \overline{D}_T(a)$ . Then,  $T|_{\overline{D}_T(a)} = T|_{\overline{D}_{T'}(a)}$ , and the claim follows immediately.

Assume that  $S$  satisfies MON and SSP. Since we have already shown that SSP is inherited, it remains to be shown that  $\mathring{S}$  satisfies MON. Let  $T = (A, \succ)$  be a tournament, and consider two alternatives  $a, b \in A$  such that  $a \in \mathring{S}(T)$  and  $b \succ a$ . Let  $T' = (A, \succ')$  be the tournament such that  $T'|_{A \setminus \{a\}} = T|_{A \setminus \{a\}}$  and  $D_{\succ'}(a) = D_{\succ}(a) \cup \{b\}$ . We have to show that  $a \in \mathring{S}(T')$ . To this end, we claim that for all  $c \in A \setminus \{a\}$ ,

$$a \notin S(\overline{D}_{\succ'}(c), \succ') \text{ implies } S(\overline{D}_{\succ}(c), \succ) = S(\overline{D}_{\succ'}(c), \succ'). \quad (1)$$

Consider the case when  $c \neq b$ . If  $a \in S(\overline{D}_{\succ}(c), \succ)$ , monotonicity of  $S$  implies that  $a \in S(\overline{D}_{\succ'}(c), \succ')$ . Equivalently,  $a \notin S(\overline{D}_{\succ'}(c), \succ')$  implies  $a \notin S(\overline{D}_{\succ}(c), \succ)$ . Now, since  $S$  satisfies SSP,

$$\begin{aligned} S(\overline{D}_{\succ'}(c), \succ') &= S(\overline{D}_{\succ'}(c) \setminus \{a\}, \succ') \text{ and} \\ S(\overline{D}_{\succ}(c), \succ) &= S(\overline{D}_{\succ}(c) \setminus \{a\}, \succ). \end{aligned}$$

It is easily verified that  $(\overline{D}_{\succ'}(c) \setminus \{a\}, \succ') = (\overline{D}_{\succ}(c) \setminus \{a\}, \succ)$ , thus we have  $S(\overline{D}_{\succ'}(c), \succ') = S(\overline{D}_{\succ}(c), \succ)$ .

If  $c = b$ , then  $a \notin S(\overline{D}_{\succ'}(b), \succ')$  together with SSP of  $S$  implies  $S(\overline{D}_{\succ'}(b), \succ') = S(\overline{D}_{\succ'}(b) \setminus \{a\}, \succ')$ . Furthermore, by definition of  $T$  and  $T'$ ,  $(\overline{D}_{\succ'}(b) \setminus \{a\}, \succ') = (\overline{D}_{\succ}(b), \succ)$  and thus  $S(\overline{D}_{\succ'}(b) \setminus \{a\}, \succ') = S(\overline{D}_{\succ}(b), \succ)$ . This proves (1).

We proceed to show that  $a \in \mathring{S}(T')$ . Assume for contradiction that this was not the case. We claim that this implies that

$$\mathring{S}(T') \text{ is } S\text{-retentive in } T. \quad (2)$$

To see this, consider  $c \in \mathring{S}(T')$ . We have to show that  $S(\overline{D}_{\succ}(c), \succ) \subseteq \mathring{S}(T')$ . Since, by assumption,  $a \notin \mathring{S}(T')$ , we have that  $a \notin S(\overline{D}_{\succ'}(c), \succ')$ . We can thus apply (1) and get

$$S(\overline{D}_{\succ}(c), \succ) = S(\overline{D}_{\succ'}(c), \succ') \text{ for all } c \in \mathring{S}(T'),$$

which, together with the  $S$ -retentiveness of  $\mathring{S}(T')$  in  $T'$ , implies (2).

Since the minimal  $S$ -retentive set is unique, it follows from (2) that  $\mathring{S}(T) \subseteq \mathring{S}(T')$ . Thus, our assumption  $a \notin \mathring{S}(T')$  implies  $a \notin \mathring{S}(T)$ , a contradiction. Hence, the assumption was false and we have  $a \in \mathring{S}(T')$ , which proves that  $\mathring{S}$  satisfies MON. This completes the proof of (ii).

For (iii), we show that the computation of  $S$  and the computation of  $\mathring{S}$  are equivalent under polynomial-time reductions.

To see that  $\mathring{S}$  can be reduced to  $S$ , consider an arbitrary tournament  $T = (A, \succ)$  and define the relation  $R = \{(a, x) : x \in S(\overline{D}(a))\}$ . It is easily verified that  $\mathring{S}(T)$  is the union of all minimal  $R$ -undominated sets or, equivalently, the maximal elements of the asymmetric part of the transitive closure of  $R$ . Observing that both  $R$  and the minimal  $R$ -undominated sets can be computed in polynomial time (see, e.g., [7], for the latter) completes the reduction.

For reduction from  $S$  to  $\mathring{S}$ , consider a tournament  $T$  on  $A$  and define  $T^* = C(T, I_a, I_b)$  for  $a, b \notin A$ . By Lemma 1,

$S(T) = \mathring{S}(T^*) \setminus \{a, b\}$ . Clearly,  $T^*$  can be computed in polynomial time from  $T$ , and  $S(T)$  can be computed in polynomial time from  $\mathring{S}(T^*)$ .  $\square$

We conclude this section by showing that, among all tournament solutions that are defined as a minimal retentive set,  $TEQ$  is the only one that satisfies COM.

**PROPOSITION 2.** *Let  $S$  be a proper tournament solution such that  $\mathcal{R}_S$  is pairwise intersecting. Then,  $\mathring{S}$  is composition-consistent if and only if  $S = TEQ$ .*

**PROOF.** It is well-known that  $TEQ$  is composition-consistent [20]. For the direction from left to right, let  $S$  be a tournament solution different from  $TEQ$ , and assume that  $\mathring{S}$  is composition-consistent. Since  $TEQ$  is the only tournament solution  $S'$  such that  $S' = \mathring{S}'$ , there has to exist a tournament  $T$  on  $A$  such that  $S(T) \neq \mathring{S}(T)$ . Let  $a, b \notin A$ , and define  $T^* = C(T, I_a, I_b)$ . By Lemma 1,  $\mathring{S}(T^*) = \{a, b\} \cup S(T)$ .

On the other hand, by composition-consistency of  $\mathring{S}$ ,  $\mathring{S}(T^*) = \mathring{S}(T) \cup \mathring{S}(I_a) \cup \mathring{S}(I_b) = \{a, b\} \cup \mathring{S}(T)$ . Hence,  $S(T) = \mathring{S}(T)$ , a contradiction.  $\square$

Although  $TRIV$  is not proper, it is easily seen that all the statements of Theorem 1 and Proposition 2 also hold for  $TRIV$ . This is due to the fact that Lemma 1 trivially holds for  $S = TRIV$ .

## 4. CONVERGENCE

In this section, we study the iterated application of the  $\circ$ -operator. Inductively define

$$S^{(0)} = S \text{ and } S^{(k+1)} = S^{\circ(k)},$$

and consider the sequence  $(S^{(n)})_{n \in \mathbb{N}} = (S^{(0)}, S^{(1)}, S^{(2)}, \dots)$ . We say that  $(S^{(n)})_{n \in \mathbb{N}}$  converges to a tournament solution  $S'$  if for each tournament  $T$ , there exists  $k_T \in \mathbb{N}$  such that  $S^{(n)}(T) = S'(T)$  for all  $n \geq k_T$ .

A perhaps surprising result is the following.

**THEOREM 2.** *Every tournament solution converges to  $TEQ$ .*

**PROOF.** Let  $S$  be a tournament solution. We show by induction on  $n$  that

$$S^{(n-1)}(T) = TEQ(T).$$

for all tournaments  $T = (A, \succ)$  of order  $|A| \leq n$ . The case  $n = 1$  is trivial. For the induction step, let  $T = (A, \succ)$  be a tournament of order  $|A| = n + 1$ . We have to show that  $S^{(n)}(T) = TEQ(T)$ . Since  $S^{(n)}$  is defined as the union of all minimal  $S^{(n-1)}$ -retentive sets, it suffices to show that a subset  $B \subseteq A$  is  $S^{(n-1)}$ -retentive if and only if it is  $TEQ$ -retentive. We have the following chain of equivalences:

$$\begin{aligned} B \text{ is } S^{(n-1)}\text{-retentive} &\text{ iff for all } b \in B, S^{(n-1)}(\overline{D}(b)) \subseteq B \\ &\text{ iff for all } b \in B, TEQ(\overline{D}(b)) \subseteq B \\ &\text{ iff } B \text{ is } TEQ\text{-retentive.} \end{aligned}$$

In particular, the second equivalence follows from the induction hypothesis, since obviously  $|\overline{D}(a)| \leq n$  for all  $a \in A$ .  $\square$

We proceed by identifying properties of  $S^{(k)}$  that are equivalent to Conjecture 1. The following lemma will be useful.

LEMMA 2. Let  $S_1$  and  $S_2$  be tournament solutions such that  $S_1 \subseteq S_2$  and  $\mathcal{R}_{S_1}$  is pairwise intersecting. Then,  $\mathcal{R}_{S_2}$  is pairwise intersecting and  $\mathring{S}_1 \subseteq \mathring{S}_2$ .

PROOF. First observe that  $S_1 \subseteq S_2$  implies that every  $S_2$ -retentive set is  $S_1$ -retentive. Now assume for contradiction that  $\mathcal{R}_{S_2}$  does not intersect pairwise and consider a tournament  $T = (A, \succ)$  with two disjoint  $S_2$ -retentive sets  $B, C \subseteq A$ . Then, by the above observation,  $B$  and  $C$  are  $S_1$ -retentive, which contradicts the fact that  $\mathcal{R}_{S_1}$  is pairwise intersecting.

Furthermore, for every tournament  $T$ ,  $\mathring{S}_2(T)$  is  $S_1$ -retentive and thus contains the unique minimal  $S_1$ -retentive set, i.e.,  $\mathring{S}_1(T) \subseteq \mathring{S}_2(T)$ .  $\square$

THEOREM 3. Let  $S$  be a tournament solution with  $TEQ \subseteq S$  that satisfies WSP or IUA. Then, the following statements are equivalent:

- (i) For all  $k \in \mathbb{N}$ ,  $\mathcal{R}_{S^{(k)}}$  is pairwise intersecting.
- (ii) For all  $k \in \mathbb{N}$ ,  $S^{(k)}$  satisfies a property  $P \in \{(\text{MON} \wedge \text{SSP}), \text{SSP}, \text{WSP}, \text{IUA}\}$  if  $S$  does.
- (iii) Conjecture 1 holds.

PROOF. To see that (i) implies (ii), assume that  $\mathcal{R}_{S^{(k)}}$  is pairwise intersecting. Then, by Theorem 1, the properties  $(\text{MON} \wedge \text{SSP})$ ,  $\text{SSP}$ ,  $\text{WSP}$ , and  $\text{IUA}$  are inherited from  $S^{(k)}$  to  $S^{(k+1)}$ .

For the implication from (ii) to (iii), let  $P \in \{\text{MON}, \text{SSP}, \text{WSP}, \text{IUA}\}$  be a basic property such that  $S^{(k)}$  satisfies  $P$  for all  $k \in \mathbb{N}$  and assume for contradiction that Conjecture 1 does not hold. We know from the work of Laffond et al. [18] and Houy [16, 17] that this assumption is equivalent to  $TEQ$  not satisfying any of the four basic properties. In particular, the latter has to be true for  $P$ . Let  $T_1$  and  $T_2$  be two tournaments showing that  $TEQ$  indeed violates  $P$ , and let  $n$  be the order of  $T_1$ . In the proof of Theorem 2, we have shown that  $S^{(n-1)}(T) = TEQ(T)$  for all tournaments  $T$  of order at most  $n$ . Thus  $T_1$  and  $T_2$  serve as an example that  $S^{(k)}$  violates  $P$ .

Finally, for the implication from (iii) to (i), assume that Conjecture 1 holds. We first prove by induction on  $k$  that  $TEQ \subseteq S^{(k)}$  for all  $k \in \mathbb{N}$ . The case  $k = 1$  holds by assumption. Now let  $T$  be a tournament and suppose that  $TEQ(T) \subseteq S^{(k)}(T)$  for some  $k \in \mathbb{N}$ . By definition,  $S^{(k+1)}(T)$  is  $S^{(k)}$ -retentive. We can thus apply the induction hypothesis to obtain that  $S^{(k+1)}(T)$  is  $TEQ$ -retentive. Since the minimal  $TEQ$ -retentive set is unique, it is contained in any  $TEQ$ -retentive set, and we have that  $TEQ(T) \subseteq S^{(k+1)}(T)$ . We can now apply Lemma 2 with  $S_1 = TEQ$  and  $S_2 = S^{(k)}$  to show that  $\mathcal{R}_{S^{(k)}}$  is pairwise intersecting for all  $k \in \mathbb{N}$ .  $\square$

Among the tournament solutions that satisfy the requirements of Theorem 3 are  $TRIV$ ,  $TC$ , the uncovered set  $UC$ , and the Banks set  $BA$  (see, e.g., [21], for definitions of the latter two).

## 4.1 Contracting Sequences

Theorem 2 showed that every tournament solution converges to  $TEQ$ . From a practical point of view, monotonic convergence that either yields smaller and smaller supersets of  $TEQ$  or larger and larger subsets of  $TEQ$  would be particularly desirable. The latter is somewhat problematic as no

refinement of  $TEQ$  is known and it is doubtful whether any such refinement would be efficiently computable. The former type of convergence turns out to be particularly useful. Call a sequence  $(S^{(n)})_{n \in \mathbb{N}}$  of tournament solutions *contracting* if for all  $k \in \mathbb{N}$ ,  $S^{(k+1)} \subseteq S^{(k)}$ . The elements of such a sequence constitute better and better “approximations” of  $TEQ$ . The following proposition identifies a sufficient condition for a sequence to be contracting.

PROPOSITION 3. Let  $S$  be a tournament solution with  $TEQ \subseteq S$ . If Conjecture 1 holds and  $\mathring{S} \subseteq S$ , then  $S^{(k+1)} \subseteq S^{(k)}$  for all  $k \in \mathbb{N}$ .

PROOF. We prove the statement by induction on  $k$ . Let  $T$  be an arbitrary tournament.  $\mathring{S}(T) \subseteq S(T)$  holds by assumption. Now suppose that  $S^{(k)}(T) \subseteq S^{(k-1)}(T)$  for some  $k \in \mathbb{N}$ . As in the proof of Theorem 3, one can show that  $TEQ \subseteq S^{(k)}$ . Applying Lemma 2 with  $S_1 = TEQ$  and  $S_2 = S^{(k)}$  yields that  $\mathcal{R}_{S^{(k)}}$  is pairwise intersecting. Therefore, we can apply Lemma 2 again, this time with  $S_1 = S^{(k)}$  and  $S_2 = S^{(k-1)}$ , which gives  $S^{(k+1)} \subseteq S^{(k)}$ .  $\square$

For example, the well-known tournament solutions  $TRIV$ ,  $TC$ ,  $UC$ , and  $MC$  give rise to contracting sequences: For  $TRIV$  and  $TC = TRIV$ , the assumptions of Proposition 3 are obviously satisfied. For  $MC$ , Laffond et al. [18] have shown that Conjecture 1 implies  $TEQ \subseteq MC$  and Brandt [4] has shown that  $\mathring{MC} \subseteq MC$ . Finally,  $TEQ \subseteq UC$  was shown by Schwartz [24] and  $\mathring{UC} \subseteq UC$  follows from Conjecture 1 and the observation that  $\mathring{UC}(T)$  is  $UC$ -retentive for all tournaments  $T$ .

The sequences  $(TRIV^{(n)})_{n \in \mathbb{N}}$  and  $(MC^{(n)})_{n \in \mathbb{N}}$  may be of particular interest. Under the assumption that Conjecture 1 holds, those sequences are contracting and all tournament solutions in those sequences satisfy all basic properties. Furthermore, by Theorem 1 (iii),  $TRIV^{(k)}$  as well as  $MC^{(k)}$  can be computed in polynomial time for any  $k \in \mathbb{N}$ . Observe that this does not imply that  $TEQ$  can be computed efficiently (due to the fact that there exists no  $k \in \mathbb{N}$  such that  $TRIV^{(k)} = TEQ$ , which follows from Proposition 4 below). In fact, Brandt et al. [8] have shown that it is NP-hard to decide whether a given alternative is in  $TEQ$ .

One might wonder if  $MC$  is contained in the sequence  $(TRIV^{(n)})_{n \in \mathbb{N}}$ . Actually, it is easy to see that this is not the case: Since  $MC$  is known to be composition-consistent (see [20]), Proposition 2 establishes that this is not the case for any  $TRIV^{(k)}$  with  $k \geq 1$ .

## 4.2 Rate of Convergence

We may ask how many iterated applications of the  $\circ$ -operator are needed until we arrive at  $TEQ$ . While we have seen that every tournament solution converges to  $TEQ$ , it turns out that no solution other than  $TEQ$  itself does so in a finite number of steps.

For a tournament solution  $S$ , let  $k_n(S)$  be the smallest  $k \in \mathbb{N}$  such that  $S^{(k)}(T) = TEQ(T)$  for all tournaments  $T$  of order at most  $n$ .

PROPOSITION 4. Let  $S \neq TEQ$  be a proper tournament solution. For each  $n \in \mathbb{N}$ ,

$$\frac{n}{2} - \frac{n_0}{2} < k_n(S) \leq n - 1,$$

where  $n_0$  is the order of the smallest tournament  $T$  with  $S(T) \neq TEQ(T)$ .

PROOF. The upper bound follows immediately from the fact that  $S^{(n-1)}(T) = TEQ(T)$  for every tournament solution  $S$  and every tournament  $T$  of order at most  $n$ . This was shown in the proof of Theorem 2.

For the lower bound, let  $S \neq TEQ$  be a tournament solution. We inductively define a family  $T_0, T_1, T_2, \dots$  of tournaments such that  $S^{(k)}(T_k) \neq TEQ(T_k)$ . Let  $T_0 = (A_0, \succ)$  be a tournament such that  $S(T_0) \neq TEQ(T_0)$ . Given  $T_{k-1} = (A_{k-1}, \succ)$ , let  $T_k = C(T_{k-1}, I_{a_k}, I_{b_k})$ , where  $a_k, b_k \notin A_{k-1}$  are two new alternatives. Observe that  $A_k = A_0 \cup \bigcup_{\ell=1}^k \{a_\ell, b_\ell\}$ .

Repeated application of Lemma 1 yields

$$\begin{aligned} S^{(k)}(T_k) &= \{a_k, b_k\} \cup S^{(k-1)}(T_{k-1}) \\ &= \{a_k, b_k\} \cup \{a_{k-1}, b_{k-1}\} \cup S^{(k-2)}(T_{k-2}) \\ &= \dots = \bigcup_{\ell=1}^k \{a_\ell, b_\ell\} \cup S(T_0). \end{aligned}$$

Since  $S(T_0) \neq TEQ(T_0)$ , we have  $S^{(k)}(T_k) \neq TEQ^{(k)}(T_k) = TEQ(T_k)$ .

We have thus shown that  $k_{n_k}(S) > k$ , where  $n_k = |A_k|$  is the order of tournament  $T_k$ . By definition of  $T_k$ ,  $n_k = n_0 + 2k$ , and therefore  $k_{n_k}(S) > k$  implies  $k_n(S) > \frac{n}{2} - \frac{n_0}{2}$ .  $\square$

As was the case for the results in Section 3, Proposition 4 also holds for  $TRIV$  even though  $TRIV$  is not a proper tournament solution. Since  $TRIV$  and  $TEQ$  differ for every tournament with two alternatives, we immediately have  $k_n(TRIV) > \frac{n}{2} - 1$ . Furthermore, Dutta [13] constructed a tournament  $T$  of order 8 for which  $TEQ(T) \neq MC(T)$ , and thus  $k_n(MC) > \frac{n}{2} - 4$ .

Interestingly, the tournaments  $T_k$  constructed in the proof of Proposition 4 show that it might be impossible to *recognize* convergence within less than  $k_n(S)$  iterations.


## 5. THE MINIMAL TC-RETENTIVE SET

As mentioned in Section 1, it is known from earlier work that Conjecture 1 is equivalent to  $TEQ$  satisfying any of the basic properties, and the attractiveness of  $TEQ$  thus hinges on the resolution of this conjecture. In Section 3 we have looked more generally at tournament solutions  $\hat{S}$ , defined as the union of all minimal  $S$ -retentive sets for arbitrary tournament solutions  $S$ . It turned out that uniqueness of minimal retentive sets again plays an important role: if  $\mathcal{R}_S$  is pairwise intersecting, then  $\hat{S}$  inherits many desirable properties from  $S$ . In this section, we prove the equivalent of Conjecture 1 for  $\mathcal{R}_{TC}$ , thus establishing  $\hat{TC}$  as an attractive new tournament solution. Note that this result is a weaker version of Conjecture 1.

**THEOREM 4.**  $\mathcal{R}_{TC}$  is pairwise intersecting.

PROOF. Consider an arbitrary tournament  $T$  on  $A$ , and assume for contradiction that  $B$  and  $C$  are two disjoint  $TC$ -retentive sets of  $T$ . Let  $b_0 \in B$  and  $c_0 \in C$ . Without loss of generality we may assume that  $c_0 \succ b_0$ . Then,  $c_0 \in \overline{D}(b_0)$  and by  $TC$ -retentiveness of  $B$  there has to be some  $b_1 \in B$  with  $b_1 \in TC(\overline{D}(b_0))$ . We claim that for each  $m \geq 1$  there must exist  $c_1, \dots, c_m \in C$  such that for all  $i$  and  $j$  with  $0 \leq i < j \leq m$ ,

- (i)  $c_{i+1} \in TC(\overline{D}(c_i))$ ,
- (ii)  $b_0 \succ c_i$  and  $c_i \succ b_1$  if and only if  $i$  is odd, and



**Figure 2: Structure of a tournament with two disjoint  $TC$ -retentive sets. A dashed edge  $(a, b)$  indicates that  $a \in TC(\overline{D}(b))$ .**

- (iii)  $c_j \succ c_i$  if and only if  $j - i$  is odd.

Before proving this claim, let us first show that it implies the statement of the theorem. For this, consider  $i$  and  $j$  with  $0 \leq i < j \leq m$ . If  $j - i$  is odd, then  $c_i \succ c_j$  by (iii). If  $j - i$  is even, then  $c_j \succ c_{j-1}$  by (i) and  $c_{j-1} \succ c_i$  by (iii). Since the dominance relation is irreflexive and anti-symmetric,  $c_i$  and  $c_j$  must be distinct alternatives in both cases. This in turn implies that the size of  $C$  is unbounded, contradicting finiteness of  $A$ . The situation is illustrated in Figure 2. We prove the claim by induction on  $m$ . First consider the case  $m = 1$ . Since  $b_1 \succ c_0$ , and by  $TC$ -retentiveness of  $C$ , there has to be some  $c_1 \in C$  with  $c_1 \in TC(\overline{D}(c_0))$  and  $c_1 \succ b_1$ . Thus (i) is satisfied. Now, by  $TC$ -retentiveness of  $B$ ,  $c_1 \notin TC(\overline{D}(b_0))$ , so it must be the case that  $c_1 \notin \overline{D}(b_0)$ , i.e.,  $b_0 \succ c_1$ . It is now easily verified that (ii) and (iii) hold as well.

Now assume that the claim is true for some  $m \geq 1$ . We show that it also holds for  $m + 1$ .

Consider the case when  $m$  is odd (the case when  $m$  is even works analogously). By  $TC$ -retentiveness of  $C$ , there has to exist some  $c_{m+1} \in C$  with  $c_{m+1} \in TC(\overline{D}(c_m))$ , which proves (i). By the induction hypothesis, and since  $m$  is odd,  $b_0 \in \overline{D}(c_m)$ . By  $TC$ -retentiveness of  $C$  and since  $c_{m+1} \in TC(\overline{D}(c_m))$ , we must have  $c_{m+1} \succ b_0$ . Moreover, since  $b_1 \in TC(b_0)$ , and by  $TC$ -retentiveness of  $B$ ,  $b_1 \succ c_{m+1}$ . This proves (ii).

For (iii), consider an arbitrary  $i$  with  $1 \leq i \leq n$ , and first assume that  $i$  is odd. If  $i = m$ , then immediately  $c_{i+1} \succ c_i$ . If  $i < m$ , then by the induction hypothesis,  $c_i \succ c_m$  and  $b_0 \succ c_m$ . Hence,  $\{c_{m+1}, c_i, b_0\} \subseteq \overline{D}(c_m)$ . Moreover, as we have already shown,  $c_{m+1} \succ b_0$ . Assuming for contradiction that  $c_i \succ c_{m+1}$ , the three alternatives  $c_{m+1}$ ,  $c_i$ , and  $b_0$  would constitute a cycle in  $\overline{D}(c_m)$ . Then, since  $c_{m+1} \in TC(\overline{D}(c_m))$ , we would have that  $b_0 \in TC(\overline{D}(c_m))$ , contradicting  $TC$ -retentiveness of  $C$ . Thus  $c_{m+1} \succ c_i$  as desired.

Now assume that  $i$  is even. By the induction hypothesis,  $c_m \succ c_i$  and  $b_1 \succ c_i$ . Assume for contradiction that  $c_{m+1} \succ c_i$  and thus  $c_{m+1} \in \overline{D}(c_i)$ . Since  $i + 1$  is odd, we already know that  $c_{m+1} \succ c_{i+1}$ . Furthermore,  $c_{i+1} \in TC(\overline{D}(c_i))$ ,

and thus  $c_{m+1} \in TC(\overline{D}(c_i))$ . However,  $b_1 \succ c_{m+1}$  and  $b_1 \in \overline{D}(c_i)$ , and thus  $b_1 \in TC(\overline{D}(c_i))$ . This contradicts  $TC$ -retentiveness of  $C$ .  $\square$

Consequently,  $\hat{TC}$  is an efficiently computable refinement of  $TC$  that satisfies all basic properties.

## 6. DISCUSSION

Assuming Schwartz's conjecture and starting with the trivial tournament solution, we have defined an infinite sequence of efficiently computable tournament solutions that are strictly contained in each other, strictly contain  $TEQ$ , and share most of its desirable properties. The implications of these findings are both of theoretical and practical nature.

From a practical point of view, we have outlined an anytime algorithm for computing  $TEQ$  that returns smaller and smaller supersets of  $TEQ$ , which are furthermore consistent according to standard properties suggested in the literature. Previous algorithms for  $TEQ$  (see, e.g., [8]) are incapable of providing *any* useful information when stopped prematurely.

From a theoretical point of view, the new perspective on  $TEQ$  as the limit of an infinite sequence of tournament solutions may prove useful for showing Schwartz's conjecture. In particular, it yields an infinite sequence of increasingly difficult conjectures, each of which is a weaker version of Schwartz's conjecture. We proved the second statement of this sequence. Our inheritance results can be interpreted as alternative proofs for the fact that Schwartz's conjecture implies that  $TEQ$  satisfies all basic properties. A natural way to prove Schwartz's conjecture would be to prove all statements of the above mentioned sequence by induction, i.e., by showing that  $\mathcal{R}_S$  is pairwise intersecting if  $\mathcal{R}_S$  is. Both proving and disproving that  $\mathcal{R}_S$  is pairwise intersecting for some reasonable solution concept  $S$  turns out to be surprisingly difficult. So far, we only have degenerate examples of tournament solutions that admit disjoint retentive sets.

## ACKNOWLEDGEMENTS

This material is based upon work supported by the Deutsche Forschungsgemeinschaft under grants BR-2312/6-1 (within the European Science Foundation's EUROCORES program LogICCC) and BR 2312/3-2.

## REFERENCES

- [1] N. Alon. Ranking tournaments. *SIAM Journal on Discrete Mathematics*, 20(1):137–142, 2006.
- [2] K. J. Arrow and H. Raynaud. *Social Choice and Multicriterion Decision-Making*. MIT Press, 1986.
- [3] D. Bouyssou, T. Marchant, M. Pirlot, A. Tsoukiàs, and P. Vincke. *Evaluation and Decision Models: Stepping Stones for the Analyst*. Springer-Verlag, 2006.
- [4] F. Brandt. Minimal stable sets in tournaments. Technical report, <http://arxiv.org/abs/0803.2138>, 2009. Presented at the 9th International Meeting of the Society of Social Choice and Welfare.
- [5] F. Brandt and F. Fischer. Computational aspects of covering in dominance graphs. In R. C. Holte and A. Howe, editors, *Proceedings of the 22nd AAAI Conference on Artificial Intelligence (AAAI)*, pages 694–699. AAAI Press, 2007.
- [6] F. Brandt and P. Harrenstein. Characterization of dominance relations in finite coalitional games. *Theory and Decision*, 2009. To appear.
- [7] F. Brandt, F. Fischer, and P. Harrenstein. The computational complexity of choice sets. In D. Samet, editor, *Proceedings of the 11th Conference on Theoretical Aspects of Rationality and Knowledge (TARK)*, pages 82–91. ACM Press, 2007.
- [8] F. Brandt, F. Fischer, P. Harrenstein, and M. Mair. A computational analysis of the tournament equilibrium set. In D. Fox and C. P. Gomes, editors, *Proceedings of the 23rd AAAI Conference on Artificial Intelligence (AAAI)*, pages 38–43. AAAI Press, 2008.
- [9] V. Conitzer. Computing Slater rankings using similarities among candidates. In *Proceedings of the 21st National Conference on Artificial Intelligence (AAAI)*, pages 613–619. AAAI Press, 2006.
- [10] J. Duggan and M. Le Breton. Dutta's minimal covering set and Shapley's saddles. *Journal of Economic Theory*, 70:257–265, 1996.
- [11] P. M. Dung. On the acceptability of arguments and its fundamental role in nonmonotonic reasoning, logic programming and n-person games. *Artificial Intelligence*, 77:321–357, 1995.
- [12] P. E. Dunne. Computational properties of argumentation systems satisfying graph-theoretic constraints. *Artificial Intelligence*, 171(10-15):701–729, 2007.
- [13] B. Dutta. On the tournament equilibrium set. *Social Choice and Welfare*, 7(4):381–383, 1990.
- [14] D. C. Fisher and J. Ryan. Tournament games and positive tournaments. *Journal of Graph Theory*, 19(2):217–236, 1995.
- [15] I. J. Good. A note on Condorcet sets. *Public Choice*, 10:97–101, 1971.
- [16] N. Houy. Still more on the tournament equilibrium set. *Social Choice and Welfare*, 32:93–99, 2009.
- [17] N. Houy. A few new results on TEQ. Unpublished Manuscript, 2009.
- [18] G. Laffond, J.-F. Laslier, and M. Le Breton. More on the tournament equilibrium set. *Mathématiques et sciences humaines*, 31(123):37–44, 1993.
- [19] G. Laffond, J.-F. Laslier, and M. Le Breton. The bipartisan set of a tournament game. *Games and Economic Behavior*, 5:182–201, 1993.
- [20] G. Laffond, J. Lainé, and J.-F. Laslier. Composition-consistent tournament solutions and social choice functions. *Social Choice and Welfare*, 13:75–93, 1996.
- [21] J.-F. Laslier. *Tournament Solutions and Majority Voting*. Springer-Verlag, 1997.
- [22] H. Moulin. Choice functions over a finite set: A summary. *Social Choice and Welfare*, 2:147–160, 1985.
- [23] H. Moulin. *Axioms of Cooperative Decision Making*. Cambridge University Press, 1988.
- [24] T. Schwartz. Cyclic tournaments and cooperative majority voting: A solution. *Social Choice and Welfare*, 7: 19–29, 1990.
- [25] G. J. Woeginger. Banks winners in tournaments are difficult to recognize. *Social Choice and Welfare*, 20: 523–528, 2003.